

SUBJECT INDEX

Actinides	2.8	- - in coupled systems	5.1, 8.4.1
ADC	7.3, 9.1	- - in dimers	5.3.1, 5.3.2
Alkyl groups	2.3	- - in heteronuclei	2.5
Amines	2.3	- - in lanthanides	2.8.3
Aminotroponoimines	2.3	- - in tetranuclear systems	5.4.2
Ammonia	2.3	Copper(II)	
Aniline	2.3	- ammonia	2.3
Aqua complexes	2.3, 3.4.1	- aqua	2.3, 3.4.1
		- -cobalt(II) dimer	5.3.3
Benzylamide	2.3	- imidazole	2.3
Benzylamine	2.3	- -nickel(II) dimer	5.3.3
Bipyridyl	2.3	Correlation function	3.2
Bloembergen equation	3.5	- time	3.2
Boltzmann population	1.5, 5.1	- - , electronic	3.2
		- - , exchange	3.2, 4.2
^{13}C	2.3, 2.5, 8.4.1	- - for contact coupling	3.2
- in amines	2.3	- - for dipolar coupling	3.2
- in cyanide	2.5	- - for electron relaxation	3.3.1
- in nickel(II) pyridine	2.3	COSY	7.1, 7.5, 9.4.2
- in nickel(II) pyridine <i>N</i> -oxide	2.3	- DQ	7.5
- in sp^2 fragments	2.5	- ISECR	7.1, 7.5
Chemical exchange	4.1	- magnitude	7.5, 9.4.2
- between two sites	4.2, 4.3	- phase sensitive	7.5, 9.4.2
- , temperature dependence	4.3.3	- ZQ	7.1, 7.5
Chemical shift	1.7.3	Cross correlation	7.8
Chloroform	1.7.3	- polarization	8.4.1
Chromium(III)		- relaxation	6.2
- aqua	2.3	Curie relaxation	3.6, 3.8, 7.8, 9.1.4
- pyridine	2.3	Cyano Met-myoglobin	2.3, 7.9
- pyridine <i>N</i> -oxide	2.3	Cytochrome c Met80Ala mutant	9.1.3
Cobalt(II)		- c'	2.3
- ammonia	2.3	- c_3	7.2, 7.4
- aqua	2.3, 3.4.1		
- -copper(II) dimer	5.3.3	δ	1.7.3
- dimer	5.3.1	DANTE	9.1.1
- imidazole	2.3	Density matrix	AV.3
- metallothioneins	5.4.2, 7.3	Diffusion	4.5.2
- -nickel(II) dimer	5.3.3	Dipolar coupling	2.2.2
- -zinc(II) dimer	5.3.1	- energy	1.1, 2.2.2, 6.2, AII
Contact relaxation	3.5, 3.8		
- shift	2.2.1, AII		

- relaxation 3.4, 3.8
- - in dimers 5.2
- - in lanthanides 3.4
- Electron relaxation 3.3
- mechanisms 3.3.1
- Enhancement factor 4.4.2
- Evans method 4.5.3
- EXSY 7.1, 7.2
- Eyring equation 4.3.3
- ^{19}F 2.5, 8.2, 8.2.2
- Ferredoxins
 - $[\text{Fe}_2\text{S}_2]^{2+}$ 5.3.2
 - $[\text{Fe}_2\text{S}_2]^+$ 5.3.4
 - $[\text{Fe}_3\text{S}_4]^+$ 5.4.1
 - $[\text{Fe}_4\text{S}_4]^{2+}$ 5.4.3, 7.2
 - $[\text{Fe}_4\text{S}_4]^{3+}$ 5.4.4, 6.2.3
 - $[\text{Fe}_4\text{S}_4]^+$ 5.4.5, 7.2
 - $[\text{Fe}_3\text{NiS}_4]^+$ 5.4.6
- FID 1.7.2, 6.1, 9.1
- Fourier transform 1.7.2, 3.2
- g factor 1.2
- g tensor 1.4
- g -anisotropy 2.2, 3.3.1, 3.7
- ^2H 2.3, 5.3.2
- relaxation 3.9
- solid state 8.3, 8.4
- Heisenberg Hamiltonian 5.1
- Hetcor 7.7, 9.5
- Heteronuclei 2.5, 7.7
- HMQC 7.7
- Hyperfine contact coupling 2.1, 2.2.1
- angular dependence 2.4
- in coupled systems 5.1
- in polymetallic systems 5.4
- in water nuclei 2.3
- Hyperfine shift 2.1
- Inversion recovery 1.7.4, 9.2
- Iron(II)
 - aqua 2.3
 - imidazole 2.3
 - intermediate spin 2.3
- Iron(III)
 - aqua 2.3, 3.4.1
 - imidazole 2.3
 - porphyrin 2.3, 2.6
- Iron in ferredoxins, dinuclear 5.3.2, 5.3.4
- , trinuclear 5.4.1
- , tetranuclear 5.4.3–5.4.5, 6.2.3, 7.2, 7.3, 7.6
- Iron sulfur proteins, high potential, oxidized 5.4.4, 6.2.3, 9.3
- - reduced 5.4.3
- Iron, trinuclear 8.3
- Jahn–Teller effect 3.3.1
- J -coupling, electronic 5.1, 8.4.1
- , nuclear 7.1, 7.5–7.7, AV
- Karplus relationship 2.4
- Lanthanides 2.8
- acetate 8.4.1
- aqua 2.3
- contact relaxation 3.5
- Curie relaxation 3.6
- dipolar relaxation 3.4
- contact shift 2.8.3
- pseudocontact shift 2.8.3
- Lattice 1.7.1
- Linewidth 1.7.4
- Magic angle sample spinning 8.1, 8.4
- Magnetic coupling 5.1
- field 1.1
- - , effective 1.7.3
- moments 1.1
- - , induced 1.6, 5.1
- - of electrons 1.2, 1.4
- - of nuclei 1.2, 1.3
- - , orbital 1.4
- susceptibility 1.6
- - tensor 2.2.2
- Magnetization 1.6
- Magnetogyric ratio 1.2
- Manganese(II)
 - ammonia 2.3
 - aqua 2.3, 3.4.1
- Metallothioneins 5.4.2, 7.3
- Met myoglobin
 - aqua 9.3
 - cyanide 2.3, 7.9
- MLEV17 7.1, 7.6
- Moment, second 8.2
- ^{14}N 2.3
- in amine 2.3
- in picoline 2.3
- in pyridine 2.3

- | | | | |
|-----------------------------|-----------------|----------------------------------|------------------------------|
| ¹⁵ N | | Radicals | 3.11, 3.12 |
| - in cyanide | 2.5 | Raman relaxation mechanism | 3.3.1 |
| - relaxation | 3.9 | R_1 | 1.7.1 |
| Neodymium(III) | 8.3 | $R_{1\rho}$ | 1.7.4 |
| Nickel(II) | | $R_{1\rho M}$ | 3.4 |
| - amine | 2.3 | R_{1e} | 3.3.1 |
| - aminotroponimine | 2.3 | R_{1M} | 3.4–3.6, 4.3.1, 5.2.1 |
| - ammonia | 2.3 | | |
| - aniline | 2.3 | R_{1p} | 4.3.2, 4.3.3 |
| - aqua | 2.3, 3.4.1 | R_2 | 1.7.1 |
| - bis(alkylxantate) | 2.3 | R_{2e} | 3.3.1 |
| - bis(salicylaldehyde) | 2.3, 7.2, 7.3 | R_{2M} | 3.4–3.6, 4.3.1, 5.2.1, 9.1.4 |
| - phenolate | 2.3 | | |
| - phosphate | 2.3 | R_{2p} | 4.3.2, 4.3.3 |
| - pyridine | 2.3 | Relaxation and magnetic coupling | 5.2 |
| - pyridine <i>N</i> -oxide | 2.3 | - and non selective experiments | 3.13, 6.2.2, 9.2 |
| - tri-iron | 5.4.6 | - and selective experiments | 3.13, 6.2.2, 9.2 |
| NOE | 6.1, 9.3, AIV | - and temperature | 3.10 |
| - in the rotating frame | 6.5 | - , Curie | 3.6, 9.1.4 |
| - measurements | 9.3 | - , electronic | 3.3 |
| - steady state | 6.2 | - in heterodimers | 5.2.2 |
| - transient | 6.4, 9.4.1 | - , ligand centered | 3.4.1 |
| - truncated | 6.3 | - , nuclear | 3.1, 3.4, 6.2 |
| NOE–NOESY | 7.9, 9.4.1 | - , outer sphere | 4.5.2 |
| NOESY | 7.1, 7.3, 9.4.1 | - , paramagnetic | 4.3.2 |
| Nuclear properties | AI | ROE | 6.5 |
| - relaxation and rotation | 3.1 | ROESY | 7.1, 7.4 |
| - - and chemical exchange | 3.1 | Rotating frame | 1.7.1 |
| - - and electron relaxation | 3.1 | | |
| - - , dipolar coupling | 3.4.1 | ¹¹⁹ Sn | 8.4.2 |
| | | $\langle S_z \rangle$ | 1.6, 2.1, 2.2.1, 5.1 |
| ¹⁷ O | 2.3, 3.4.1 | Salicylaldehydes | 2.3 |
| Orbach relaxation mechanism | 3.3.1 | Saturation transfer | 4.3.4, 7.2 |
| Orbital magnetic moment | 1.4 | Shielding constant | 1.7.3 |
| Outer sphere relaxation | 4.5.2 | Shift | |
| | | - bulk susceptibility | 4.5.3 |
| Phenanthroline | 2.3 | - , contact | 2.2.1, 2.5, 5.1, AII |
| Phenylisonitrile | 2.3 | | |
| Phosphine | 2.3 | - , pseudocontact | 2.2.2, AIII |
| Phosphine <i>N</i> -oxide | 2.3 | - reagents | 2.8.5 |
| Picoline | 2.3 | Simulation of spectra | AV |
| Porphyrines | 2.3 | SOD | |
| - , tetraphenyl | 2.3 | - , CoCo | 5.3.3 |
| ppm | 1.7.3 | - , CuCo | 5.3.3, 6.2.3 |
| Precession | 1.2 | - , CuCu | 5.2.2 |
| Presaturation | 9.1.1 | - , CuNi | 5.3.3 |
| Product operators | AV.2 | - , CuZn | 5.2.1 |
| Protoporphyrin IX | 2.3 | - , ECo | 5.3.3, 9.3 |
| Pseudocontact shifts | 2.2.2 | Solomon equation | 3.4 |
| - , factorization | 2.7, 2.8.4 | Spectral density | 3.2 |
| - in high spin iron(III) | 2.6 | Spin density | 2.1 |
| - in lanthanides | 2.8.2 | - diffusion | 6.2.3 |
| - in low spin iron(III) | 2.3 | - eigenvalues | 1.2 |
| - in metalloproteins | 2.9 | | |

- | | | | |
|-----------------------------|-------------------|---------------------------|----------|
| - Hamiltonian | 1.4 | Uranium trinuclear | 8.4.1 |
| - lattice relaxation | 1.7.1 | | |
| - moments | 1.2 | Vanadium(III) aqua | 2.3 |
| - orbit | 1.4 | Vanadyl(IV) aqua | 2.3 |
| - polarization | 2.1 | Van Vleck equation | 1.6 |
| - rotation | 3.3.1 | Viscosity | 3.2, 3.6 |
| - σ -spin relaxation | 1.7.1 | | |
| - wavefunctions | 1.2 | | |
| Superweft | 9.1.3 | Water complexes | 2.3 |
| | | Watergate | 9.1.2 |
| τ_c | 3.2 | Weft | 9.1.3 |
| τ_M | 3.2, 4.2, 4.3.3 | | |
| τ_r | 3.2 | ^{89}Y | 8.4.2 |
| τ_s | 3.2, 5.2.2 | Ytterbium porphinate | 7.5 |
| τ_v | 3.3.1 | | |
| T_1 | 1.7.1, 6.2.2, 9.2 | Zeeman operator | |
| - , non-selective | 3.13, 6.2.2, 9.2 | - , electronic | 1.4 |
| - , selective | 3.13, 6.2.2, 9.2 | - , nuclear | 1.3 |
| Tempol | 8.4 | ZFS | 1.4, 2.3 |
| Titanium(III) aqua | 2.3 | - and electron relaxation | 3.3.1 |
| TMS | 1.7.3 | - and nuclear relaxation | 3.7 |
| TOCSY | 7.1, 7.6 | | |
| - , clean | 9.4.2 | | |

